

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

factors. Environmental factors were found to have a marked influence upon invasion by the parasite (Fusarium), there being a "critical soil temperature" (about 17° C.) for such invasion. Below this the plants are not invaded even in the sickest soils.—J. M. C.

Desiccation.—An investigation of the course of desiccation and partial starvation in cacti has been made by MacDougal, Long, and Brown.<sup>33</sup> The principal studies center upon the changing rate of water loss, chemical changes in the food reserves, plasmatic colloids and cell sap, and the morphological changes which occur during long periods of desiccation. In one case a large *Echinocactus* was under observation for 6 years after removal of the plant from the soil. Water loss is rather rapid at first, but proceeds more and more slowly with time. While 10 per cent of the water was lost the first year in one specimen, during the sixth year only 5 per cent of the water remaining at the beginning of that year was lost. The loss of water is much more rapid of course in the open than in diffuse light, and *Echinocactus* can withstand desiccation not more than 2 years with free exposure.—Geo. D. Fuller.

Aeration of nutrient solutions.—STILES and JÖRGENSEN<sup>34</sup> find that aeration of the nutrient solution increases the rate of growth of barley, as found by various workers, but has no effect on the growth of buckwheat, as found by FREE. They carefully limit their conclusion to the condition under which they experimented, and find themselves unable to explain this specific difference. They emphasize the necessity of knowing much more about the physical chemistry of water culture solutions. They also feel that neither the law of the minimum nor the principle of limiting factors gives an adequate expression of the behavior of the plant as a whole.—WM. CROCKER.

Apogamy in ferns.—Steil.<sup>35</sup> has discovered apogamy in a large number of ferns, the investigation extending over a period of 6 years. It seems that apogamy is of frequent occurrence in *Pellaea*, *Pteris*, and *Aspidium*. The prothallia were grown under cultural conditions favorable for the development of sex organs and embryos in non-apogamous species. Many interesting details of embryo development are given, which much extend our knowledge of this phenomenon.—J. M. C.

<sup>&</sup>lt;sup>33</sup> MACDOUGAL, D. T., LONG, E. R., and BROWN, J. G., End results of desiccation and respiration in succulent plants. Physiol. Res. r:289-325. 1915.

<sup>&</sup>lt;sup>34</sup> STILES, W., and JÖRGENSEN, I., Observations on the influence of aeration of the nutrient solution in water culture experiments, with some remarks on the water culture methods. New Phytol. 16:182-197. 1917.

<sup>&</sup>lt;sup>35</sup> STEIL, W. N., Studies of some new cases of apogamy in ferns. Bull. Torr. Bot. Club 45:93-108. *pls.* 4, 5. 1918.